

ON THE EXISTENCE OF SOLUTIONS TO THE OPERATOR RICCATI EQUATION AND THE $\tan \Theta$ THEOREM

V. KOSTRYKIN, K. A. MAKAROV, AND A. K. MOTOVILOV

ABSTRACT. Let A and C be self-adjoint operators such that the spectrum of A lies in a gap of the spectrum of C and let $d > 0$ be the distance between the spectra of A and C . We prove that under these assumptions the sharp value of the constant c in the condition $\|B\| < cd$ guaranteeing the existence of a (bounded) solution to the operator Riccati equation $XA - CX + XBX = B^*$ is equal to $\sqrt{2}$. We also prove an extension of the Davis-Kahan $\tan \Theta$ theorem and provide a sharp estimate for the norm of the solution to the Riccati equation. If C is bounded, we prove, in addition, that the solution X is a strict contraction if B satisfies the condition $\|B\| < d$, and that this condition is sharp.

1. INTRODUCTION

In this paper we consider the operator Riccati equation

$$XA - CX + XBX = B^* \quad (1.1)$$

associated with the self-adjoint 2×2 block operator matrix

$$H = \begin{pmatrix} A & B \\ B^* & C \end{pmatrix} \quad (1.2)$$

on the orthogonal sum $\mathcal{H} = \mathcal{H}_A \oplus \mathcal{H}_C$ of separable Hilbert spaces $\mathcal{H}_A$ and $\mathcal{H}_C$. Here A is a bounded self-adjoint operator on the Hilbert space $\mathcal{H}_A$, C possibly unbounded self-adjoint operator on the Hilbert space $\mathcal{H}_C$, and B is a bounded operator from $\mathcal{H}_C$ to $\mathcal{H}_A$.

Solving the Riccati equation appears to be an adequate tool in the study of the invariant subspaces of the operator H that are the graphs of bounded operators from $\mathcal{H}_A$ to $\mathcal{H}_C$, the so-called graph subspaces. It is well known that given a bounded solution $X : \mathcal{H}_A \rightarrow \mathcal{H}_C$ to the Riccati equation (1.1) (with $\text{Ran } X \subset \text{Dom}(C)$), the graph

$$\mathcal{G}(X) = \{x \oplus Xx \mid x \in \mathcal{H}_A\}$$

of the operator X reduces the operator H (see, e.g., [3, Section 5]) and in the framework of this approach the following two problems naturally arise. The first problem is to study the spectrum of the part of H associated with the reducing subspace $\mathcal{G}(X)$ (respectively its orthogonal complement $\mathcal{G}(X)^\perp$), and the second one is to estimate the operator angle Θ (see, e.g., [10] for discussion of this notion) between the subspaces $\mathcal{G}(X)$ and $\mathcal{H}_A$ (respectively $\mathcal{G}(X)^\perp$ and $\mathcal{H}_C$).

2000 *Mathematics Subject Classification.* Primary 47A55, 47A15; Secondary 47B15.

Key words and phrases. invariant subspaces, operator Riccati equation, operator matrix, operator angle, factorization theorem.

Both of these problems can efficiently be solved if a bounded solution X to (1.1) is known: the operator H appears to be similar to the diagonal block operator matrix

$$\begin{pmatrix} A + BX & 0 \\ 0 & C - B^*X^* \end{pmatrix}$$

associated with the decomposition $\mathcal{H} = \mathcal{H}_A \oplus \mathcal{H}_C$ (in particular, $\sigma(H) = \sigma(A + BX) \cup \sigma(C - B^*X^*)$), and the operator angle Θ between the subspaces $\mathcal{G}(X)$ and $\mathcal{H}_A$ has the representation

$$\Theta = \arctan \sqrt{X^*X}.$$

If the spectra of A and C overlap, the Riccati equation (1.1) may have no solution at all (cf., e.g., [3, Example 3.2]). At the same time the spectra separation requirement alone does not guarantee the existence of solutions either (see, e.g., [3, Lemma 3.11]). Under the spectra separation hypothesis

$$\text{dist}(\sigma(A), \sigma(C)) > 0, \quad (1.3)$$

a natural sufficient condition for the existence of solutions to the Riccati equation (1.1) requires a smallness assumption on the operator B of the form

$$\|B\| < c_{\text{best}} \text{dist}(\sigma(A), \sigma(C)) \quad (1.4)$$

with a constant $c_{\text{best}} > 0$ independent of the distance between the spectra $\sigma(A)$ and $\sigma(C)$ of the operators A and C , respectively. The best possible constant c_{best} in (1.4) is still unknown. However, c_{best} is known to be in the interval $[\pi^{-1}, \sqrt{2}]$ (see [3]). If both A and C are bounded, then $c_{\text{best}} \in [c_\pi, \sqrt{2}]$ with $c_\pi = \frac{3\pi - \sqrt{\pi^2 + 32}}{\pi^2 - 4} = 0.503288\dots$ (see [12]). In [12] the best possible constant c_{best} has been conjectured to be $\sqrt{3}/2$. Some earlier results in this direction can be found in [2], [17], [19], and [20].

In some particular cases the optimal solvability condition (1.4) can be relaxed provided that some additional assumptions upon mutual disposition of the spectra of A and C are posed. For instance, if the spectra of A and C are subordinated, e.g.,

$$\sup \sigma(A) < \inf \sigma(C), \quad (1.5)$$

the Riccati equation (1.1) is known to have a strictly contractive solution for any bounded B (see, e.g., [1]). To some extent abusing the terminology one may say that in this case the best possible constant in inequality (1.4) is infinite: No smallness assumptions on B are needed.

In the limiting case of (1.5),

$$\sup \sigma(A) = \inf \sigma(C),$$

the existence of contractive solutions has been established in [2] under some additional assumptions which have been dropped in [11]. See also [18] where the spectra separation condition (1.3) has also been somewhat relaxed and the existence of a bounded but not necessarily contractive solution has been established.

Our *first principal result* concerns the case where the operator C has a finite spectral gap containing the spectrum of A . Recall that by a finite spectral gap of a self-adjoint operator T one understands an *open* finite interval on the real axis lying in the resolvent set of T such that both of its end points belong to the spectrum of T .

Theorem 1. Assume that the self-adjoint operator C has a finite spectral gap Δ containing the spectrum of the bounded self-adjoint operator A .

(i) Suppose that

$$\|B\| < \sqrt{d|\Delta|} \quad \text{where} \quad d = \text{dist}(\sigma(A), \sigma(C)), \quad (1.6)$$

with $|\cdot|$ denoting Lebesgue measure on $\mathbb{R}$. Then the spectrum of the block operator matrix $H = \begin{pmatrix} A & B \\ B^* & C \end{pmatrix}$ in the gap Δ is a (proper) closed subset of (the open set) Δ . The spectral subspace of the operator H associated with the interval Δ is the graph of a bounded solution $X : \mathcal{H}_A \rightarrow \mathcal{H}_C$ to the Riccati equation (1.1). Moreover, the operator X is the unique solution to the Riccati equation in the class of bounded operators with the properties

$$\sigma(A + BX) \subset \Delta \quad \text{and} \quad \sigma(C - B^*X^*) \subset \mathbb{R} \setminus \Delta. \quad (1.7)$$

(ii) If in addition the operator C is bounded and

$$\|B\| < \sqrt{d(|\Delta| - d)}, \quad (1.8)$$

then the solution X is a strict contraction and

$$\|X\| \leq \tan \frac{1}{2} \arctan \left(\frac{2\|AB + BC\|}{d(|\Delta| - d) - \|B\|^2} \right) < 1. \quad (1.9)$$

As a corollary, under the assumption that the operator C has a finite spectral gap Δ containing the spectrum of A , we prove that $c = \sqrt{2}$ is best possible in condition (1.4) ensuring the existence of a bounded solution to the Riccati equation (1.1) (see Remark 3.3) while $c = 1$ is best possible in (1.4) to ensure that the solution is a contraction (see Remark 5.3).

The proof of the part (i) of Theorem 1 will be given in Section 3 and that of the part (ii) in Section 5.

Our *second principal result* holds with no *a priori* assumption upon the mutual disposition of the spectra of A and C (in particular, the spectra of A and C may overlap).

Theorem 2. Assume that the self-adjoint operator C has a spectral gap Δ (finite or infinite) and the self-adjoint operator A is bounded. Assume that the Riccati equation (1.1) has a bounded solution X and hence the graph subspace $\mathcal{G}(X)$ reduces the block operator matrix H . Suppose that the spectrum of the part $H|_{\mathcal{G}(X)}$ of the operator H associated with the reducing subspace $\mathcal{G}(X)$ is a (proper) closed subset of (the open set) Δ . Then the following norm estimate holds:

$$\|X\| \leq \frac{\|B\|}{\delta} \quad \text{with} \quad \delta = \text{dist}(\sigma(H|_{\mathcal{G}(X)}), \sigma(C)). \quad (1.10)$$

Equivalently,

$$\|\tan \Theta\| \leq \frac{\|B\|}{\delta}, \quad (1.11)$$

where Θ is the operator angle between the subspaces $\mathcal{H}_A$ and $\mathcal{G}(X)$.

Estimate (1.11) extends the Davis-Kahan $\tan \Theta$ theorem [7], a result previously known only in the case where the spectra of C and $H|_{\mathcal{G}(X)}$ are subordinated, that is, the operator C is semi-bounded and the spectrum of the part $H|_{\mathcal{G}(X)}$ lies in the *infinite* spectral gap of C . This generalization extends the list of the celebrated Davis-Kahan $\sin \Theta$ and $\sin 2\Theta$ theorems, proven in the case where the operator C has a gap of finite length [7].

The proof of Theorem 2 will be given in Section 4.

Our main techniques are based on applications of the Virozub-Matsaev factorization theorem for analytic operator-valued functions [22] (in the spirit of the work [18] (cf. [15])) and the Daletsky-Krein factorization formula [5]. Under the hypothesis of Theorem 1 we prove that

- for $\lambda \notin \sigma(C)$ the operator-valued Herglotz function $M(\lambda) = \lambda I - A + B(C - \lambda I)^{-1}B^*$ admits a factorization

$$M(\lambda) = W(\lambda)(Z - \lambda I), \quad (1.12)$$

with W being an operator-valued function holomorphic on the resolvent set of the operator C and Z is a bounded operator with the spectrum in the spectral gap Δ of the operator C ,

- the Riccati equation (1.1) has a bounded solution of the form

$$X = -\frac{1}{2\pi i} \int_{\Gamma} d\lambda (C - \lambda I)^{-1} B^* (Z - \lambda I)^{-1}, \quad (1.13)$$

where Γ is an appropriate Jordan contour encircling the spectrum of the operator Z ,

- the spectral subspace of the 2×2 block operator matrix H (1.2) associated with the interval Δ is the graph of the operator X ,
- the spectrum of the operator H in the interval Δ coincides with that of the operator Z , that is, $\sigma(H) \cap \Delta = \sigma(Z)$.

In Section 2 we recall the concept of invariant graph subspaces for linear operators as well as their relation to the Riccati equation. Theorem 2.5 below presents a general result linking the factorization property (1.12) of the operator valued function $M(\lambda)$ with the existence of a spectral subspace for the 2×2 self-adjoint block operator matrix (1.2) admitting representation as the graph of the operator (1.13). In Section 3 under hypothesis (1.6) we prove factorization formula (1.12) and give bounds on the location of the spectrum of the operator Z (Theorem 3.2), and finally prove the part (i) of Theorem 1. The proof of Theorem 2 is given in Section 4. In Section 5 combining the results of Theorems 1 (i) and 2 under assumptions (1.6) and (1.8) respectively we provide norm estimates on the solution X of the Riccati equation and prove Theorem 1 (ii). We conclude the section by an example showing that condition (1.8) ensuring the strict contractivity of the solution X is sharp.

Few words about notations used throughout the paper. Given a Hilbert space $\mathcal{K}$ by $I_{\mathcal{K}}$ we denote the identity operator on $\mathcal{K}$. If it does not lead to any confusion we will simply write I instead of more pedantic notation $I_{\mathcal{K}}$. The set of all bounded linear operators from the Hilbert space $\mathcal{K}$ to a Hilbert space $\mathcal{L}$ will be denoted by $\mathcal{B}(\mathcal{K}, \mathcal{L})$. If $\mathcal{L} = \mathcal{K}$ the shorthand $\mathcal{B}(\mathcal{K})$ will be used for this set. Let K and L be self-adjoint operators on a Hilbert space $\mathcal{K}$. We say $K < L$ (or, equivalently, $L > K$) if there is a number $\gamma > 0$ such that $L - K > \gamma I$. The notation $\rho(T)$ will be used for the resolvent set of a closed operator T .

After completing this work we learned that a result similar to the part (ii) of Theorem 1 has been recently obtained within a different approach by A. V. Selin (private communication).

2. INVARIANT GRAPH SUBSPACES AND BLOCK DIAGONALIZATION

In this section we collect some results related to the invariant graph subspaces of a linear operator as well as to the closely related problem of block diagonalization of block operator matrices.

First, recall the definition a graph subspace.

Definition 2.1. Let $\mathcal{K}$ be a closed subspace of a Hilbert space $\mathcal{N}$ and $X \in \mathcal{B}(\mathcal{K}, \mathcal{K}^\perp)$. Denote by $P_{\mathcal{K}}$ and $P_{\mathcal{K}^\perp}$ the orthogonal projections in $\mathcal{N}$ onto the subspace $\mathcal{K}$ and orthogonal complement $\mathcal{K}^\perp$, respectively. The set

$$\mathcal{G}(X) = \{x \in \mathcal{N} \mid P_{\mathcal{K}^\perp}x = XP_{\mathcal{K}}x\}$$

is called the graph subspace associated with the operator X .

For notational setup we assume the following

Hypothesis 2.2. Let H_0 be a self-adjoint operator in a Hilbert space $\mathcal{H}$ and $\mathcal{H}_A \subset \mathcal{H}$ a reducing subspace of H_0 . Assume that with respect to the decomposition

$$\mathcal{H} = \mathcal{H}_A \oplus \mathcal{H}_C \quad (\mathcal{H}_C = \mathcal{H} \ominus \mathcal{H}_A) \quad (2.1)$$

the operator H_0 reads as the block diagonal operator matrix

$$H_0 = \text{diag}(A, C)$$

with A being a bounded self-adjoint operator in $\mathcal{H}_A$, C a possibly unbounded self-adjoint operator in $\mathcal{H}_C$, and $\text{Dom}(H_0) = \mathcal{H}_A \oplus \text{Dom}(C)$. Assume, in addition, that with respect to the decomposition (2.1) the self-adjoint operator H reads as

$$H = \begin{pmatrix} A & B \\ B^* & C \end{pmatrix}, \quad \text{Dom}(H) = \mathcal{H}_A \oplus \text{Dom}(C), \quad (2.2)$$

where B is a bounded operator from $\mathcal{H}_C$ to $\mathcal{H}_A$.

Under Hypothesis 2.2, a bounded operator X from $\mathcal{H}_A$ to $\mathcal{H}_C$ is said to be a solution to the Riccati equation (1.1) if $\text{Ran}(X) \subset \text{Dom}(C)$ and (1.1) holds as an operator equality.

The existence of a bounded solution to the Riccati equation (1.1) is equivalent to a possibility of the block diagonalization of the operator matrix H with respect to the decomposition $\mathcal{H} = \mathcal{G}(X) \oplus \mathcal{G}(X)^\perp$. The precise statement is as follows (see Lemma 5.3 and Theorem 5.5 in [3]; also cf. [2], [6], [10], [20]).

Theorem 2.3. Assume Hypothesis 2.2. Then a bounded operator X from $\mathcal{H}_A$ to $\mathcal{H}_C$ is a solution to the Riccati equation (1.1) iff the graph $\mathcal{G}(X)$ of X reduces the 2×2 block operator matrix H . Moreover, if $X \in \mathcal{B}(\mathcal{H}_A, \mathcal{H}_C)$ is a solution to (1.1) then:

(i) The operator $V^{-1}HV$ with

$$V = \begin{pmatrix} I & -X^* \\ X & I \end{pmatrix}$$

is block diagonal with respect to decomposition (2.1). Furthermore,

$$V^{-1}HV = \begin{pmatrix} Z & 0 \\ 0 & \widehat{Z} \end{pmatrix},$$

where $Z = A + BX$ with $\text{Dom}(Z) = \mathcal{H}_A$ and $\widehat{Z} = C - B^*X^*$ with $\text{Dom}(\widehat{Z}) = \text{Dom}(C)$.

(ii) The operator

$$\Lambda = (I + X^*X)^{1/2}Z(I + X^*X)^{-1/2} \quad (2.3)$$

and possibly unbounded operator

$$\widehat{\Lambda} = (I + XX^*)^{1/2}\widehat{Z}(I + XX^*)^{-1/2} \quad (2.4)$$

with $\text{Dom}(\widehat{\Lambda}) = (I + XX^*)^{1/2}(\text{Dom}(C))$ are self-adjoint operators in $\mathcal{H}_A$ and $\mathcal{H}_C$, respectively.

Theorem 2.3 yields the following uniqueness result as a corollary.

Corollary 2.4. *Assume Hypothesis 2.2. Suppose that Σ and $\widehat{\Sigma}$ are disjoint Borel subsets of $\mathbb{R}$ such that $\text{dist}(\Sigma, \widehat{\Sigma}) > 0$. Let $\mathcal{X} = \mathcal{X}(A, B, \Sigma, \widehat{\Sigma})$ be the set of all bounded operators X from $\mathcal{H}_A$ to $\mathcal{H}_C$ with the properties*

$$\sigma(A + BX) \subset \Sigma, \quad (2.5)$$

$$\sigma(C - B^*X^*) \subset \widehat{\Sigma}, \quad \text{Dom}(C - B^*X^*) = \text{Dom}(C). \quad (2.6)$$

Then if $X, Y \in \mathcal{X}$ satisfy the Riccati equation (1.1), then $X = Y$.

Proof. Suppose X and Y are two bounded solutions to (1.1) both satisfying (2.5) and (2.6). Then by Theorem 2.3 the graphs of X and Y both coincide with the spectral subspace of the 2×2 operator matrix (2.2) associated with the set Σ , and hence, $X = Y$. $\square$

Under Hypothesis 2.2 introduce the operator-valued Herglotz function

$$M(\lambda) = \lambda I - A + B(C - \lambda I)^{-1}B^*, \quad \lambda \in \rho(C). \quad (2.7)$$

By definition the spectrum $\sigma(M)$ of the function M is the set of all $\lambda \in \mathbb{C}$ such that either the operator $M(\lambda)$ is not invertible or the inverse $[M(\lambda)]^{-1}$ is an unbounded operator.

It is well known (see, e.g., [17]) that the resolvent of the operator H (2.2) can be represented as the following 2×2 operator matrix

$$\begin{aligned} (H - \lambda I)^{-1} &= \begin{pmatrix} 0 & 0 \\ 0 & (C - \lambda I)^{-1} \end{pmatrix} \\ &\quad - \begin{pmatrix} I \\ -(C - \lambda I)^{-1}B^* \end{pmatrix} M(\lambda)^{-1} \begin{pmatrix} I & -B(C - \lambda I)^{-1} \end{pmatrix}, \\ &\quad \lambda \in \rho(H), \end{aligned} \quad (2.8)$$

where M is the Herglotz function given by (2.7). Representation (2.8) shows that for $\lambda \in \rho(C)$ the operator $H - \lambda I$ has a bounded inverse iff $M(\lambda)$ does, which means that

$$\sigma(H) \cap \rho(C) = \sigma(M) \cap \rho(C). \quad (2.9)$$

We will also need the following general result (cf. [2, Theorem 2.2], [18, Proposition 2.4 and Theorem 2.5], and [15, Theorems 4.4 and 5.1]).

Theorem 2.5. *Assume Hypothesis 2.2 and suppose that the Herglotz function (2.7) admits the factorization*

$$M(\lambda) = W(\lambda)(Z - \lambda I), \quad \lambda \in \Omega, \quad (2.10)$$

where Z is a bounded operator in $\mathcal{H}_A$ such that $\sigma(Z) \cap \sigma(C) = \emptyset$, Ω is a domain in $\rho(C)$ such that $\sigma(Z) \subset \Omega$, and W is a holomorphic $\mathcal{B}(\mathcal{H}_A)$ -valued function on Ω , such that for any $\lambda \in \sigma(Z)$

the operator $W(\lambda)$ has a bounded inverse. Then $\sigma(Z)$ is an isolated part of the spectrum of the operator H and the spectral subspace $\text{Ran } E_H(\sigma(Z))$ of H associated with the set $\sigma(Z)$ is the graph of the bounded operator X from $\mathcal{H}_A$ to $\mathcal{H}_C$ given by

$$X = -\frac{1}{2\pi i} \int_{\Gamma} d\lambda (C - \lambda I)^{-1} B^* (Z - \lambda I)^{-1}. \quad (2.11)$$

Here, Γ is an arbitrary Jordan contour in $\rho(Z) \cap \rho(C)$ (maybe consisting of several simple Jordan contours) encircling $\sigma(Z)$ in the clockwise direction and having winding number 0 with respect to the spectrum of C .

Moreover, the operator Z can be written in terms of the operator X given by (2.11) as

$$Z = A + BX \quad (2.12)$$

and the factor $W(\lambda)$ admits analytic continuation to the whole resolvent set of the operator C by the following formula

$$W(\lambda) = I - B(C - \lambda)^{-1}X, \quad \lambda \in \rho(C). \quad (2.13)$$

Proof. By hypothesis the function $W(\lambda)$ is holomorphic on an open set $\Omega \subset \rho(C)$ containing the closed subset $\sigma(Z)$ and the operator $W(\lambda)$ has a bounded inverse for any $\lambda \in \sigma(Z)$. Hence there is an open neighborhood $\tilde{\Omega}$ of $\sigma(Z)$ in Ω where the operator $W(\lambda)$ is boundedly invertible, i.e. $W(\lambda)^{-1} \in \mathcal{B}(\mathcal{H}_A)$ for any $\lambda \in \tilde{\Omega}$. By (2.10)

$$M(\lambda)^{-1} = (Z - \lambda I)^{-1} W(\lambda)^{-1}, \quad \lambda \in \tilde{\Omega} \setminus \sigma(Z). \quad (2.14)$$

Taking into account (2.9) one infers that the spectrum of H in $\tilde{\Omega}$ coincides with that of M and, thus, with that of Z , that is,

$$\sigma(H) \cap \tilde{\Omega} = \sigma(Z), \quad (2.15)$$

Since $\tilde{\Omega}$ is an open set, $\sigma(Z)$ is a closed set, and $\sigma(Z) \subset \tilde{\Omega}$, one also concludes that $\sigma(Z)$ is isolated from the remaining part of the spectrum of H , i.e.

$$\text{dist}(\sigma(Z), \sigma(H) \setminus \sigma(Z)) > 0.$$

Using representation (2.8), for the spectral projection $E_H(\sigma(Z))$ of the operator H associated with the set $\sigma(Z)$ the Riesz integration yields

$$\begin{aligned} E_H(\sigma(Z)) &= E_H(\sigma(Z)) = \frac{1}{2\pi i} \int_{\Gamma} d\lambda (H - \lambda I)^{-1} \\ &= -\frac{1}{2\pi i} \int_{\Gamma} d\lambda \begin{pmatrix} I \\ -(C - \lambda I)^{-1} B^* \end{pmatrix} M(\lambda)^{-1} \begin{pmatrix} I & -B(C - \lambda I)^{-1} \end{pmatrix}, \end{aligned}$$

where Γ stands for an arbitrary Jordan contour (possibly consisting of several simple Jordan contours) in $\tilde{\Omega}$ encircling the spectrum of Z in the clockwise direction and having winding number 0 with respect to the spectrum of C . Hence,

$$E_H(\sigma(Z)) = \begin{pmatrix} E & G^* \\ G & F \end{pmatrix}, \quad (2.16)$$

where

$$\begin{aligned} E &= -\frac{1}{2\pi i} \int_{\Gamma} d\lambda M(\lambda)^{-1}, \\ F &= -\frac{1}{2\pi i} \int_{\Gamma} d\lambda (C - \lambda I)^{-1} B^* M(\lambda)^{-1} B (C - \lambda I)^{-1}, \end{aligned} \quad (2.17)$$

and

$$\begin{aligned} G &= \frac{1}{2\pi i} \int_{\Gamma} d\lambda (C - \lambda I)^{-1} B^* M(\lambda)^{-1} \\ &= \frac{1}{2\pi i} \int_{\Gamma} d\lambda (C - \lambda I)^{-1} B^* (Z - \lambda I)^{-1} W(\lambda)^{-1}, \end{aligned}$$

using factorization formula (2.10). Since both the operator-valued functions $(C - \lambda)^{-1}$ and $W(\lambda)^{-1}$ are holomorphic in $\tilde{\Omega}$, applying the Daletsky-Krein formula (see [5, Lemma I.2.1]) we get

$$\begin{aligned} G &= \left[\frac{1}{2\pi i} \int_{\Gamma} d\lambda (C - \lambda I)^{-1} B^* (Z - \lambda I)^{-1} \right] \\ &\quad \times \left[\frac{1}{2\pi i} \int_{\Gamma} d\lambda (Z - \lambda I)^{-1} W(\lambda)^{-1} \right]. \end{aligned} \quad (2.18)$$

Hence, combining (2.14) and (2.17) proves the representation

$$G = XE, \quad (2.19)$$

where X (the first factor on the r.h.s. part of (2.18)) is given by

$$X = -\frac{1}{2\pi i} \int_{\Gamma} d\lambda (C - \lambda I)^{-1} B^* (Z - \lambda I)^{-1}. \quad (2.20)$$

In an analogous way one also proves that

$$F = XEX^*. \quad (2.21)$$

Clearly, for $\lambda \in \Gamma$ we have $\text{Ran}(C - \lambda I)^{-1} \subset \text{Dom}(C)$ and hence

$$\text{Ran}(X) \subset \text{Dom}(C),$$

which immediately follows from (2.20). Multiplying both sides of (2.20) by B from the left yields

$$BX = -\frac{1}{2\pi i} \int_{\Gamma} d\lambda B(C - \lambda I)^{-1} B^* (Z - \lambda I)^{-1}. \quad (2.22)$$

Meanwhile,

$$B(C - \lambda I)^{-1} B^* = A - \lambda I + M(\lambda) = A - \lambda I + W(\lambda)(Z - \lambda I), \quad \lambda \in \Gamma,$$

and, hence, using (2.22)

$$BX = -\frac{1}{2\pi i} \int_{\Gamma} d\lambda [W(\lambda) + A(Z - \lambda I)^{-1} - \lambda(Z - \lambda I)^{-1}]. \quad (2.23)$$

The function $W(\lambda)$ is holomorphic in the domain bounded by the contour Γ and, thus, the first term in the integrand on the r.h.s. of (2.23) gives no contribution. Since Γ encircles the spectrum

of Z , the integration of the remaining two terms in (2.23) can be performed explicitly using the operator version of the residue theorem, which yields $BX = -A + Z$ and hence

$$Z = A + BX \quad (2.24)$$

proving (2.12).

Since the spectra of the operators C and Z are disjoint and Z is a bounded operator, it is straightforward to show (see, e.g., [4] or [21]) that the operator X given by (2.20) is the unique solution to the operator Sylvester equation

$$XZ - CX = B^*,$$

which by (2.24) proves that X solves the Riccati equation (1.1).

Now applying Theorem 2.3 and using (2.16), (2.19), and (2.21) we arrive at the series of equalities

$$\begin{aligned} I &= \frac{1}{2\pi i} \int_{\Gamma} d\lambda (Z - \lambda I)^{-1} \\ &= \frac{1}{2\pi i} \int_{\Gamma} d\lambda P_A \begin{pmatrix} (Z - \lambda I)^{-1} & 0 \\ 0 & (\widehat{Z} - \lambda I)^{-1} \end{pmatrix} P_A^* \\ &= \frac{1}{2\pi i} \int_{\Gamma} d\lambda P_A V^{-1} (H - \lambda I)^{-1} V P_A^* \\ &= P_A V^{-1} E_H(\sigma(Z)) V P_A^* \\ &= E(I + X^* X), \end{aligned} \quad (2.25)$$

where $\widehat{Z} = C - B^* X^*$ with $\text{Dom}(\widehat{Z}) = \text{Dom}(C)$,

$$V = \begin{pmatrix} I & -X^* \\ X & I \end{pmatrix},$$

and P_A is the canonical projection from $\mathcal{H}$ to $\mathcal{H}_A$ (i.e., $P_A(f_A \oplus f_C) = f_A$ for $f_A \in \mathcal{H}_A$ and $f_C \in \mathcal{H}_C$). Combining (2.19), and (2.25) one concludes that the spectral projection $E_H(\sigma(Z))$ admits the representation

$$E_H(\sigma(Z)) = \begin{pmatrix} (I + X^* X)^{-1} & (I + X^* X)^{-1} X^* \\ X(I + X^* X)^{-1} & X(I + X^* X)^{-1} X^* \end{pmatrix}. \quad (2.26)$$

Note that the contour Γ in (2.20) can be replaced by an arbitrary Jordan contour in $\rho(Z) \cap \rho(C)$ (possibly consisting of several simple Jordan contours) encircling the set $\sigma(Z)$ in the clockwise direction and having winding number 0 with respect to $\sigma(C)$. Then observing that the r.h.s. of (2.26) is nothing but the orthogonal projection in $\mathcal{H} = \mathcal{H}_A \oplus \mathcal{H}_C$ onto the graph of the operator X proves that $\text{Ran } E_H(\sigma(Z)) = \mathcal{G}(X)$.

We conclude with the proof of the representation (2.13). First, by (2.24) we notice that

$$\begin{aligned} [I - B(C - \lambda)^{-1} X](Z - \lambda) &= [I - B(C - \lambda)^{-1} X](A + BX - \lambda) \\ &= A + BX - \lambda - B(C - \lambda)^{-1}(XA + XBX - \lambda X), \\ &\quad \lambda \in \rho(C). \end{aligned}$$

Since X solves the Riccati equation (1.1) one infers that $XA + XBX = B^* + CX$ which implies

$$\begin{aligned} [I - B(C - \lambda)^{-1}X](Z - \lambda) &= M(\lambda) + BX - B(C - \lambda)^{-1}(C - \lambda)X \\ &= M(\lambda), \quad \lambda \in \rho(C). \end{aligned} \quad (2.27)$$

Hence combining (2.10) and (2.27) yields

$$W(\lambda) = M(\lambda)(Z - \lambda)^{-1} = I - B(C - \lambda)^{-1}X, \quad \lambda \in \Omega \setminus \sigma(Z).$$

Then the analytic continuation completes the proof. $\square$

Remark 2.6. By analytic continuation of both parts of (2.10) one concludes that the factorization formula (2.10) holds for any $\lambda \in \rho(C)$ with $W(\lambda)$ given by (2.13). Moreover, the representation

$$[W(\lambda)]^{-1} = (Z - \lambda I)[M(\lambda)]^{-1}, \quad \lambda \in \tilde{\Omega} \setminus \sigma(M) = \tilde{\Omega} \setminus \sigma(Z),$$

and the fact that $\sigma(M) \subset \sigma(H)$ imply that $[W(\lambda)]^{-1}$ admits analytic continuation as a $\mathcal{B}(\mathcal{H}_A)$ -valued holomorphic function to the domain $\rho(H) \cup \sigma(Z)$.

3. AN EXISTENCE RESULT. PROOF OF THEOREM 1 (I)

As we have already mentioned in Introduction our main technical tool in proving the solvability of the Riccati equation is the Virozub-Matsaev factorization theorem [22] (also see [16]). For convenience of the reader we reproduce the corresponding statement following Propositions 1.1 and 1.2 in [18].

Theorem 3.1. *Let $\mathcal{X}$ be a Hilbert space and $F(\lambda)$ a holomorphic $\mathcal{B}(\mathcal{X})$ -valued function on a simply connected domain $\Omega \subset \mathbb{C}$. Assume that Ω includes an interval $[a, b] \subset \mathbb{R}$ such that*

$$F(a) < 0, \quad F(b) > 0, \quad \text{and} \quad \frac{d}{d\lambda}F(\lambda) > 0 \quad \text{for all} \quad \lambda \in [a, b].$$

Then there exist a domain $\tilde{\Omega} \subset \Omega$ containing $[a, b]$ and a unique bounded operator Z on $\mathcal{X}$ with $\sigma(Z) \subset (a, b)$ such that $F(\lambda)$ admits the factorization

$$F(\lambda) = G(\lambda)(Z - \lambda I), \quad \lambda \in \tilde{\Omega},$$

where $G(\lambda)$ is a holomorphic operator-valued function on $\tilde{\Omega}$ whose values are bounded and boundedly invertible operators in $\mathcal{X}$, that is,

$$G(\lambda) \in \mathcal{B}(\mathcal{X}) \quad \text{and} \quad [G(\lambda)]^{-1} \in \mathcal{B}(\mathcal{X}) \quad \lambda \in \tilde{\Omega}.$$

Based on Theorem 3.1 we obtain the following factorization result, the cornerstone for our further considerations.

Theorem 3.2. *Assume Hypothesis 2.2. Assume, in addition, that C has a finite spectral gap $\Delta = (\alpha, \beta)$, $\alpha < \beta$, the spectrum of A lies in Δ , i.e., $\sigma(A) \subset \Delta$, and*

$$\|B\| < \sqrt{d|\Delta|},$$

where

$$d = \text{dist}(\sigma(A), \sigma(C)).$$

Then there is a unique operator $Z \in \mathcal{B}(\mathcal{H}_A)$ with $\sigma(Z) \subset \Delta$ such that the operator-valued function $M(\lambda)$ given by (2.7) admits the factorization

$$M(\lambda) = W(\lambda)(Z - \lambda I), \quad \lambda \in \rho(C), \quad (3.1)$$

with a holomorphic $\mathcal{B}(\mathcal{H}_A)$ -valued function W on $\rho(C)$. Moreover, for any

$$\lambda \in (\mathbb{C} \setminus \sigma(M)) \cup \Delta$$

the operator $W(\lambda)$ has a bounded inverse and

$$\sigma(Z) = \sigma(H) \cap \Delta \subset [\inf \sigma(A) - \delta_-, \sup \sigma(A) + \delta_+], \quad (3.2)$$

where

$$\delta_- = \|B\| \tan \left(\frac{1}{2} \arctan \frac{2\|B\|}{\beta - \inf \sigma(A)} \right) < \inf \sigma(A) - \alpha, \quad (3.3)$$

$$\delta_+ = \|B\| \tan \left(\frac{1}{2} \arctan \frac{2\|B\|}{\sup \sigma(A) - \alpha} \right) < \beta - \sup \sigma(A). \quad (3.4)$$

Proof. By the spectral theorem

$$B(C - \lambda I)^{-1}B^* = \int_{\mathbb{R} \setminus \Delta} B E_C(d\mu) B^* \frac{1}{(\mu - \lambda)}, \quad \lambda \in \rho(C),$$

where $E_C(\mu)$ stands for the spectral family of the self-adjoint operator C . Hence

$$\frac{d}{d\lambda} M(\lambda) = I + \int_{\mathbb{R} \setminus \Delta} B E_C(d\mu) B^* \frac{1}{(\mu - \lambda)^2}, \quad \lambda \in \rho(C). \quad (3.5)$$

For $\lambda \in \Delta$ the integral in (3.5) is a non-negative operator. Therefore, the derivative of $M(\lambda)$ is a strictly positive operator:

$$\frac{d}{d\lambda} M(\lambda) \geq I > 0, \quad \lambda \in \Delta.$$

Next we estimate the quadratic form of $M(\lambda)$. Let $f \in \mathcal{H}_A$, $\|f\| = 1$. Then

$$\begin{aligned} \langle M(\lambda) f, f \rangle &= \lambda - \langle A f, f \rangle + \langle (C - \lambda)^{-1} B^* f, B^* f \rangle \\ &= \lambda - \langle A f, f \rangle + \int_{-\infty}^{\alpha} \frac{1}{\mu - \lambda} \langle E_C(d\mu) B^* f, B^* f \rangle \\ &\quad + \int_{\beta}^{+\infty} \frac{1}{\mu - \lambda} \langle E_C(d\mu) B^* f, B^* f \rangle, \quad \lambda \in \rho(C). \end{aligned} \quad (3.6)$$

Since for $\lambda \in \Delta$ the integral in the second line of (3.6) is non-positive and the one in the third line is non-negative, one obtains the two-sided estimate

$$\left(\lambda - \sup \sigma(A) - \frac{\|B\|^2}{\lambda - \alpha} \right) I \leq M(\lambda) \leq \left(\lambda - \inf \sigma(A) - \frac{\|B\|^2}{\lambda - \beta} \right) I, \quad \lambda \in \Delta.$$

Now, a simple calculation shows that

$$M(\lambda) < 0 \quad \text{for} \quad \lambda \in (\alpha, \inf \sigma(A) - \delta_-) \quad (3.7)$$

and

$$M(\lambda) > 0 \quad \text{for} \quad \lambda \in (\sup \sigma(A) + \delta_+, \beta), \quad (3.8)$$

where δ_- and δ_+ are given by (3.3) and (3.4), respectively.

Thus, the function $F(\lambda) = M(\lambda)$ satisfies assumptions of Theorem 3.1 for any $a \in (\alpha, \inf \sigma(A) - \delta_-)$ and any $b \in (\sup \sigma(A) + \delta_+, \beta)$, proving the existence of the unique bounded operator $Z \in \mathcal{B}(\mathcal{H}_A)$ such that (3.1) and (3.2) hold taking into account (2.9). It follows from Theorem 3.1 that the factor $W(\lambda)$ in (2.10) has a bounded inverse in a complex neighborhood $U \subset \mathbb{C}$ of the interval $[\inf \sigma(A) - \delta_-, \sup \sigma(A) + \delta_+]$. Moreover, the operator $W(\lambda)$ has a bounded inverse for any $\lambda \in (\mathbb{C} \setminus \sigma(M)) \cup \Delta$ by Remark 2.6. The proof is complete. $\square$

We are ready to prove the part (i) of Theorem 1.

Proof of Theorem 1 (i). By Theorem 3.2 the Herglotz function (2.7) admits the factorization (2.10) with $W(\lambda)$ and Z satisfying hypothesis of Theorem 2.5. Therefore, the Riccati equation (1.1) has a bounded solution X given by (2.11). Theorem 2.5 also shows that the graph $\mathcal{G}(X)$ of the operator X coincides with the spectral subspace $\text{Ran } E_H(\sigma(Z))$ for the block operator matrix H given by (2.2). By (3.2) the subspace $\text{Ran } E_H(\sigma(Z))$ coincides with $\text{Ran } E_H(\Delta)$, proving that

$$\text{Ran } E_H(\Delta) = \mathcal{G}(X). \quad (3.9)$$

To prove that X possesses the properties (1.7) we proceed as follows. Let

$$V = \begin{pmatrix} I & -X^* \\ X & I \end{pmatrix}, \quad S = (I + X^*X)^{1/2}, \quad \text{and} \quad \widehat{S} = (I + XX^*)^{1/2}. \quad (3.10)$$

From Theorem 2.3 it follows that

$$E_H(\Delta) = V \begin{pmatrix} S^{-1}E_\Lambda(\Delta)S & 0 \\ 0 & \widehat{S}^{-1}E_{\widehat{\Lambda}}(\Delta)\widehat{S} \end{pmatrix} V^{-1}, \quad (3.11)$$

where Λ and $\widehat{\Lambda}$ are the self-adjoint operators defined by (2.3) and (2.4), respectively; $E_\Lambda(\Delta)$ and $E_{\widehat{\Lambda}}(\Delta)$ denote the spectral projections for Λ and $\widehat{\Lambda}$ associated with the interval Δ . Since Λ is similar to Z ($\Lambda = SZS^{-1}$) and by Theorem 3.2 the inclusion $\sigma(Z) \subset \Delta$ holds, one concludes that $\sigma(\Lambda) \subset \Delta$. Hence, $E_\Lambda(\Delta) = I$ and then (3.11) implies the equality

$$E_H(\Delta) = V \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} V^{-1} + V \begin{pmatrix} 0 & 0 \\ 0 & \widehat{S}^{-1}E_{\widehat{\Lambda}}(\Delta)\widehat{S} \end{pmatrix} V^{-1}. \quad (3.12)$$

The first summand on the r.h.s. of (3.12)

$$V \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} V^{-1} = \begin{pmatrix} (I + X^*X)^{-1} & (I + X^*X)^{-1}X^* \\ X(I + X^*X)^{-1} & X(I + X^*X)^{-1}X^* \end{pmatrix}$$

coincides with the orthogonal projection onto the graph subspace $\mathcal{G}(X)$ and, hence, by (3.9) it equals $E_H(\Delta)$. Thus, the second summand on the r.h.s. of (3.12) vanishes,

$$V \begin{pmatrix} 0 & 0 \\ 0 & \widehat{S}^{-1}E_{\widehat{\Lambda}}(\Delta)\widehat{S} \end{pmatrix} V^{-1} = 0,$$

which means that $E_{\widehat{\Lambda}}(\Delta) = 0$ and, therefore, $\sigma(\widehat{\Lambda}) \cap \Delta = \emptyset$. By (2.4) it is now straightforward to see that X satisfies (1.7). It remains to conclude that by Corollary 2.4 the operator X is the unique bounded solution to (1.1) satisfying (1.7) which completes the proof of Theorem 1 (i). $\square$

Remark 3.3. Obviously, under hypothesis of Theorem 1 the inequality $|\Delta| \geq 2d$ holds, with the equality sign occurring only if the spectrum of the operator A is a one point set. Hence, the condition $\|B\| < \sqrt{2}d$, which is stronger than (1.6), implies the existence of a bounded solution to the Riccati equation (1.1). This means that in the case where C has a finite spectral gap Δ and $\sigma(A) \subset \Delta$ the best possible constant c_{best} in condition (1.4) ensuring the solvability of the Riccati equation satisfies the inequality $c_{\text{best}} \geq \sqrt{2}$. On the other hand in [3, Lemma 3.11 and Remark 3.12] it is shown that $c_{\text{best}} \leq \sqrt{2}$. Thus, $c = \sqrt{2}$ is best possible in inequality (1.4) which ensures the solvability of the Riccati equation under the additional hypothesis that $\text{dist}(\sigma(A), \sigma(C)) > 0$ and that the spectrum of A lies in a spectral gap of the operator C .

4. THE TAN Θ THEOREM

We start out by recalling a concept of the operator angle between two subspaces in a Hilbert space going back to the works by Friedrichs [8], M. Krein, Kransnoselsky, and Milman [13], [14], Halmos [9], and Davis and Kahan [7]. A comprehensive discussion of this concept can be found, e.g., in [10].

Given a closed subspace Q of the Hilbert space $\mathcal{H} = \mathcal{H}_A \oplus \mathcal{H}_C$, introduce the operator angle Θ between the subspaces $\mathcal{H}_A \oplus \{0\}$ and Q by

$$\Theta = \arcsin \sqrt{I_{\mathcal{H}_A} - P_A Q P_A^*}, \quad (4.1)$$

where P_A is the canonical projection from $\mathcal{H}$ onto $\mathcal{H}_A$ and Q the orthogonal projection in $\mathcal{H}$ onto Q . If the subspace Q is the graph $\mathcal{G}(X)$ of a bounded operator X from $\mathcal{H}_A$ to $\mathcal{H}_C$, then (see [10]; cf. [7] and [9])

$$\tan \Theta = \sqrt{X^* X} \quad (4.2)$$

and

$$\|\sin \Theta\| = \|Q - P\|, \quad (4.3)$$

where $P = P_A^* P_A$ denotes the orthogonal projection in $\mathcal{H}$ onto the subspace $\mathcal{H}_A \oplus \{0\}$.

Note that the common definition of the operator angle (see, e.g., [10]) slightly differs from (4.1). Usually, the operator angle is defined as the restriction of the operator (4.1) onto the maximal subspace of $\mathcal{H}_A$ where it has a trivial kernel. Clearly, the difference in these two definitions does not effect the value of the norm $\|\tan \Theta\|$.

Now we are ready to prove the second principal result of the paper, a generalization of the Davis-Kahan tan Θ Theorem [7].

Proof of Theorem 2. By hypothesis the Riccati equation (1.1) has a bounded solution X . Then, by Theorem 2.3 the operator $Z = A + BX$ is similar to the bounded self-adjoint operator Λ given by (2.3) and hence

$$\sigma(Z) = \sigma(\Lambda) \subset \mathbb{R}. \quad (4.4)$$

The Riccati equation (1.1) can be rewritten in the form

$$X(Z - \gamma I) - (C - \gamma I)X = B^*, \quad (4.5)$$

where

$$\gamma = \frac{1}{2}(\sup \sigma(Z) + \inf \sigma(Z)) = \frac{1}{2}(\sup \sigma(\Lambda) + \inf \sigma(\Lambda)). \quad (4.6)$$

By hypothesis $\text{dist}(\sigma(H|_{\mathcal{G}(X)}), \sigma(C)) = \delta > 0$ and then Theorem 2.3 implies that

$$\sigma(H|_{\mathcal{G}(X)}) = \sigma(Z) = \sigma(\Lambda),$$

which by hypothesis proves the inclusion

$$\sigma(C) \subset (-\infty, \inf \sigma(\Lambda) - \delta] \cup [\sup \sigma(\Lambda) + \delta, \infty). \quad (4.7)$$

Hence, combining (4.4), (4.6), and (4.7) proves that $\gamma \in \rho(C)$ and

$$\|(C - \gamma I)^{-1}\| = \frac{1}{\|\Lambda - \gamma I\| + \delta}. \quad (4.8)$$

Multiplying both sides of (4.5) by $(C - \gamma I)^{-1}$ from the left one gets the representation

$$X = (C - \gamma I)^{-1}(X(Z - \gamma I) - B^*). \quad (4.9)$$

Using Theorem 2.3 (ii) one obtains the estimate

$$\begin{aligned} \|X(Z - \gamma I)\| &= \|X(I + X^*X)^{-1/2}(\Lambda - \gamma I)(I + X^*X)^{1/2}\| \\ &\leq \|X(I + X^*X)^{-1/2}\| \|\Lambda - \gamma I\| (1 + \|X\|^2)^{1/2}. \end{aligned} \quad (4.10)$$

Clearly, by the spectral theorem

$$\|X(I + X^*X)^{-1/2}\| = \sqrt{\|X^*X(I + X^*X)^{-1}\|} = \frac{\|X\|}{(1 + \|X\|^2)^{1/2}}.$$

Hence (4.10) implies the estimate

$$\|X(Z - \gamma I)\| \leq \|X\| \|\Lambda - \gamma I\|,$$

which together with (4.9) proves the norm inequality

$$\|X\| \leq \|(C - \gamma I)^{-1}\| (\|\Lambda - \gamma I\| \|X\| + \|B\|). \quad (4.11)$$

Solving inequality (4.11) with respect to $\|X\|$ and taking into account (4.8) proves the norm estimate (1.10). Finally, since $\|\sqrt{X^*X}\| = \|X\|$, by the definition of the operator angle (4.2) one gets

$$\|\tan \Theta\| = \|X\|.$$

Hence, (1.10) is equivalent to (1.11). $\square$

Remark 4.1. It is natural to ask whether estimate (1.11) remains to hold if one replaces the distance $\delta = \text{dist}(\sigma(H|_{\mathcal{G}(X)}), \sigma(C))$ by

$$\widehat{\delta} = \text{dist}(\sigma(H|_{\mathcal{G}(X)^\perp}, \sigma(A)) = \text{dist}(\sigma(C - B^*X^*), \sigma(A)).$$

The answer is “No”: Example 6.1 in [7] shows that the inequality $\widehat{\delta} \|\tan \Theta\| \leq \|B\|$ fails to hold in general.

5. NORM ESTIMATES OF SOLUTIONS. PROOF OF THEOREM 1 (II)

The existence result of Theorem 1 (i) by itself gives no clue for estimating the norm of the corresponding solution X to the Riccati equation. To the contrary, Theorem 2 provides such an estimate whenever some additional information on the spectrum location of the “perturbed” operator matrix $H = \begin{pmatrix} A & B \\ B^* & C \end{pmatrix}$ is available. In turn, the bounds on the spectrum of the part $H|_{\mathcal{G}(X)}$ of the operator matrix H associated with the reducing subspace $\mathcal{G}(X)$ (needed to satisfy the hypotheses of Theorem 2) can be obtained by combining the results of Theorems 2.5 and 3.2. As a result of performing this program one gets an *a priori* estimate on the norm of the solution X .

Theorem 5.1. *Assume hypothesis of Theorem 1 (i) with $\Delta = (\alpha, \beta)$, $\alpha < \beta$. Let X be the unique solution to the Riccati equation (1.1) referred to in Theorem 1. Then*

$$\|X\| \leq \frac{\|B\|}{\tilde{\delta}}, \quad (5.1)$$

where

$$\tilde{\delta} = \min\{\inf \sigma(A) - \alpha - \delta_-, \beta - \sup \sigma(A) - \delta_+\} > 0 \quad (5.2)$$

with $\delta_{\pm}$ given by (3.3) and (3.4).

Proof. Under hypothesis of Theorem 1 (i) with $\Delta = (\alpha, \beta)$, $\alpha < \beta$ one can apply Theorem 3.2 and using the same strategy of proof as that of Theorem 1 (i) one concludes that

$$Z = A + BX,$$

where Z is the unique operator with $\sigma(Z) \subset \Delta$ referred to in Theorem 3.2 and X is the unique solution to the Riccati equation (1.1) referred to in Theorem 1 (i). Hence, by Theorem 3.2

$$\sigma(A + BX) \subset [\inf \sigma(A) - \delta_-, \sup \sigma(A) + \delta_+]. \quad (5.3)$$

By hypothesis (of Theorem 1 (i))

$$\sigma(C) \subset (-\infty, \alpha] \cup [\beta, \infty), \quad (5.4)$$

which together with (5.3) yields the inequality

$$\text{dist}(\sigma(A + BX), \sigma(C)) \geq \tilde{\delta}.$$

By Theorem 2.5 one observes that $\sigma(H)|_{\mathcal{G}(X)} = \sigma(Z)$ which proves (5.1) using Theorem 2. $\square$

The proof of Theorem 1 (ii) needs complementary considerations. Our reasoning is based on the celebrated Davis-Kahan $\tan 2\Theta$ -Theorem [7]. For convenience of the reader we reproduce the corresponding result (cf. Theorem 2.4 (iii) and Remark 2.8 in [11] and Corollary 6.4 in [10]).

Theorem 5.2. *Assume Hypothesis 2.2. Suppose that the operator C is bounded and $\sup \sigma(A) < \inf \sigma(C)$. Then the open interval $(\sup \sigma(A), \inf \sigma(C))$ is a spectral gap of the operator H and the spectral subspace $E_H((-\infty, \sup \sigma(A)))$ is the graph of a contractive operator X from $\mathcal{H}_A$ to*

$\mathcal{H}_C$. Moreover, the operator X is the unique contractive solution to the Riccati equation (1.1) and its norm satisfies the estimate

$$\|X\| \leq \tan \left(\frac{1}{2} \arctan \frac{2\|B\|}{d} \right) < 1,$$

where

$$d = \text{dist}(\sigma(A), \sigma(C)).$$

Now we are prepared to prove Theorem 1 (ii).

Proof of Theorem 1 (ii). Let X be the solution to the Riccati equation (1.1) referred to in Theorem 1 (i) and thus the spectral subspace of the operator H associated with the interval Δ is the graph $\mathcal{G}(X)$ of the operator X . Then $\mathcal{G}(X)$ is also the spectral subspace of the operator $(H - \gamma I)^2$ associated with the interval $[0, |\Delta|^2/4]$ where γ is the center of the interval Δ , that is,

$$\mathcal{G}(X) = \text{Ran}(E_H(\Delta)) = \text{Ran}(E_{(H-\gamma I)^2}[0, |\Delta|^2/4]). \quad (5.5)$$

By inspection one obtains that with respect to the decomposition $\mathcal{H} = \mathcal{H}_A \oplus \mathcal{H}_C$ the non-negative operator $(H - \gamma I)^2$ reads

$$(H - \gamma I)^2 = \begin{pmatrix} \hat{A} & \hat{B} \\ \hat{B}^* & \hat{C} \end{pmatrix}, \quad (5.6)$$

where $\hat{A} = (A - \gamma I)^2 + BB^*$, $\hat{B} = AB + BC$, and $\hat{C} = (C - \gamma I)^2 + B^*B$.

The hypothesis that the spectrum of C lies in $\mathbb{R} \setminus \Delta$ implies the operator inequality

$$\hat{C} \geq \frac{|\Delta|^2}{4} I + BB^* \geq \frac{|\Delta|^2}{4} I.$$

The hypothesis $\sigma(A) \subset \Delta$ yields

$$0 \leq \hat{A} \leq \left(\frac{|\Delta|}{2} - d \right)^2 I + BB^* \leq \left[\left(\frac{|\Delta|}{2} - d \right)^2 + \|B\|^2 \right] I, \quad (5.7)$$

taking into account that $\text{dist}(\sigma(A), \sigma(C)) = d$. Hence, under hypothesis (1.8) one concludes that

$$\text{dist}(\sigma(\hat{A}), \sigma(\hat{C})) \geq d(|\Delta| - d) - \|B\|^2 > 0 \quad (5.8)$$

and that the spectra $\sigma(\hat{A})$ and $\sigma(\hat{C})$ of the entries $\hat{A}$ and $\hat{C}$ are subordinated, that is, $\sup \sigma(\hat{A}) < \inf \sigma(\hat{C})$. By Theorem 5.2 (cf. Theorem 2.1 in [1]) one infers that the interval $(\sup \sigma(\hat{A}), \inf \sigma(\hat{C}))$ lies in the resolvent set of the operator $(H - \gamma I)^2$. In particular, the following inclusion holds

$$((|\Delta|/2 - d)^2 + \|B\|^2, |\Delta|^2/4) \subset \rho((H - \gamma I)^2). \quad (5.9)$$

Therefore, the spectral subspaces of the operator $(H - \gamma I)^2$ associated with the intervals $[0, |\Delta|^2/4]$ and $[0, (|\Delta|/2 - d)^2 + \|B\|^2]$, respectively, coincide, that is,

$$\begin{aligned} \mathcal{G}(X) &= \text{Ran}(E_{(H-\gamma I)^2}([0, |\Delta|^2/4])) \\ &= \text{Ran}(E_{(H-\gamma I)^2}([0, (|\Delta|/2 - d)^2 + \|B\|^2])). \end{aligned} \quad (5.10)$$

From (5.6) one concludes that the operator matrix $(H - \gamma I)^2$ is an off-diagonal perturbation of the matrix $\text{diag}\{\hat{A}, \hat{C}\}$ diagonal with respect to the decomposition $\mathcal{H} = \mathcal{H}_A \oplus \mathcal{H}_C$. Applying again Theorem 5.2 proves that the spectral subspace of the operator $(H - \gamma I)^2$ associated with

the interval $[0, |(\Delta/2 - d)^2 + \|B\|^2]$ is the graph of a contraction $\widehat{X}$ satisfying the norm-estimate (1.9). By (5.5) and (5.10) this subspace coincides with $\mathcal{G}(X)$ and, therefore, $\widehat{X} = X$ and hence (1.9) holds for X , completing the proof. $\square$

Remark 5.3. Condition (1.8) ensuring the strict contractivity of the solution X is sharp. This can be seen as follows. Let $\mathcal{H}_A = \mathbb{C}$, $\mathcal{H}_C = \mathbb{C}^2$, $A = 0$,

$$C = \begin{pmatrix} -d & 0 \\ 0 & d \end{pmatrix}, \quad d > 0,$$

and

$$B = \left(\frac{b}{\sqrt{2}}, \frac{b}{\sqrt{2}} \right), \quad b \in \mathbb{R}.$$

By inspection one proves that the 2×1 matrix

$$X = \begin{pmatrix} -\frac{b}{\sqrt{2}d} \\ \frac{b}{\sqrt{2}d} \end{pmatrix} \tag{5.11}$$

solves the Riccati equation

$$XA - CX + XBX = B^*.$$

Moreover,

$$A + BX = 0$$

and X possesses the properties (1.7). Clearly, $\|B\| = b$ and $\|X\| = \frac{b}{d} = \frac{\|B\|}{d}$.

For $b \in [d, \sqrt{2}d]$ hypothesis (1.6) is satisfied with $\Delta = (-d, d)$, condition (1.8) fails to hold, and $\|X\| \geq 1$, that is, estimate (1.8) is sharp.

REFERENCES

- [1] V. M. Adamjan and H. Langer, *Spectral properties of a class of rational operator valued functions*, J. Operator Theory **33** (1995), 259 – 277.
- [2] V. Adamjan, H. Langer, and C. Tretter, *Existence and uniqueness of contractive solutions of some Riccati equations*, J. Funct. Anal. **179** (2001), 448 – 473.
- [3] S. Albeverio, K. A. Makarov, and A. K. Motovilov, *Graph subspaces and the spectral shift function*, Canad. J. Math. (to appear); arXiv: math.SP/0105142.
- [4] Yu. L. Daleckiĭ, *On the asymptotic solution of a vector differential equation*, Dokl. Akad. Nauk SSSR **92** (1953), 881 – 884 (Russian).
- [5] Ju. L. Daleckiĭ and M. G. Kreĭn, *Stability of Solutions of Differential Equations in Banach Spaces*, Translations of Mathematical Monographs, Vol. 43, AMS, Providence, Rhode Island, 1974.
- [6] J. Daughtry, *Isolated solutions of quadratic matrix equations*, Linear Algebra Appl. **21** (1978), 89 – 94.
- [7] C. Davis and W. M. Kahan, *The rotation of eigenvectors by a perturbation. III*, SIAM J. Numer. Anal. **7** (1970), 1 – 46.
- [8] K. Friedrichs, *On certain inequalities and characteristic value problems for analytic functions and for functions of two variables*, Trans. Amer. Math. Soc. **41** (1937), 321 – 364.
- [9] P. R. Halmos, *Two subspaces*, Trans. Amer. Math. Soc. **144** (1969), 381–389.
- [10] V. Kostykin, K. A. Makarov, and A. K. Motovilov, *Existence and uniqueness of solutions to the operator Riccati equation. A geometric approach*, in Yu. Karpeshina, G. Stolz, R. Weikard, Y. Zeng (Eds.), *Advances in Differential Equations and Mathematical Physics*, Contemporary Mathematics **327**, Amer. Math. Soc., 2003 (to appear); arXiv: math.SP/0207125.
- [11] V. Kostykin, K. A. Makarov, and A. K. Motovilov, *A generalization of the tan 2Θ theorem*, arXiv: math.SP/0302020.

- [12] V. Kostykin, K. A. Makarov, and A. K. Motovilov, *Perturbation of spectra and spectral subspaces*, preprint (2003).
- [13] M. G. Krein and M. A. Krasnoselsky, *Fundamental theorems about extensions of Hermite operators and some applications to the theory of orthogonal polynomials and to the moment problem*, Uspekhi Mat. Nauk **2** (1947), 60 – 106 (Russian).
- [14] M. G. Krein, M. A. Krasnoselsky, and D. P. Milman, *On defect numbers of linear operators in Banach space and some geometric problems*, Sbornik Trudov Instituta Matematiki Akademii Nauk Ukrainskoy SSR, **11** (1948), 97 – 112 (Russian).
- [15] H. Langer, A. Markus, V. Matsaev, and C. Tretter, *A new concept for block operator matrices: the quadratic numerical range*, Linear Algebra Appl. **330** (2001), 89 – 112.
- [16] A. S. Markus and V. I. Matsaev, *Spectral theory of holomorphic operator-functions in Hilbert space*, Funct. Anal. Appl. **9** (1975), 73 – 74.
- [17] R. Mennicken and A. K. Motovilov, *Operator interpretation of resonances arising in spectral problems for 2×2 operator matrices*, Math. Nachr. **201** (1999), 117 – 181; arXiv: funct-an/9708001.
- [18] R. Mennicken and A. A. Shkalikov, *Spectral decomposition of symmetric operator matrices*, Math. Nachr. **179** (1996), 259 – 273.
- [19] A. K. Motovilov, *Potentials appearing after the removal of energy-dependence and scattering by them*, in Proc. Intern. Workshop “Mathematical Aspects of the Scattering Theory and Applications”, St. Petersburg State University, St. Petersburg, 1991, pp. 101 – 108.
- [20] A. K. Motovilov, *Removal of the resolvent-like energy dependence from interactions and invariant subspaces of a total Hamiltonian*, J. Math. Phys. **36** (1995), 6647 – 6664; arXiv: funct-an/9606002.
- [21] M. Rosenblum, *On the operator equation $BX - XA = Q$* , Duke Math. J. **23** (1956), 263 – 269.
- [22] A. I. Virozub and V. I. Matsaev, *The spectral properties of a certain class of self-adjoint operator functions*, Funct. Anal. Appl. **8** (1974), 1 – 9.

Acknowledgments. V. Kostykin is grateful to V. Enss, A. Knauf, H. Leschke, and R. Schrader for useful discussions. K. A. Makarov is indebted to Graduiertenkolleg “Hierarchie und Symmetrie in mathematischen Modellen” for kind hospitality during his stay at RWTH Aachen in summer 2002. A. K. Motovilov acknowledges the kind hospitality and support by the Department of Mathematics, University of Missouri, Columbia, MO, USA. He was also supported in part by the Russian Foundation for Basic Research.

FRAUNHOFER-INSTITUT FÜR LASERTECHNIK, STEINBACHSTRASSE 15, AACHEN, D-52074, GERMANY
E-mail address: kostykin@ilt.fraunhofer.de, kostykin@t-online.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA
E-mail address: makarov@math.missouri.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA.
Permanent address: BLTP, JINR, 141980 DUBNA, MOSCOW REGION, RUSSIA
E-mail address: motovilv@thsun1.jinr.ru